

Superaligned Fermi Beta Decay

J.C. Hardy^a and I. S. Towner^b

^a*Cyclotron Institute, Texas A & M University,
College Station, TX 77843*

^b*Physics department, Queen's University,
Kingston, Ontario K7L 3N6, Canada*

Abstract. Superaligned $0^+ \rightarrow 0^+$ nuclear beta decay provides a direct measure of the weak vector coupling constant, G_V . We survey current world data on the nine accurately determined transitions of this type, which range from the decay of ^{10}C to that of ^{54}Co , and demonstrate that the results confirm conservation of the weak vector current (CVC) but differ at the 98% confidence level from the unitarity condition for the Cabibbo-Kobayashi-Maskawa (CKM) matrix. We examine the reliability of the small calculated corrections that have been applied to the data, and assess the likelihood of even higher quality nuclear data becoming available to confirm or deny the discrepancy. Some of the required experiments depend upon the availability of intense radioactive beams. Others are possible today.

CURRENT STATUS OF WORLD DATA

Superaligned Fermi $0^+ \rightarrow 0^+$ nuclear beta decays [1,2] provide both the best test of the Conserved Vector Current (CVC) hypothesis in weak interactions and, together with the muon lifetime, the most accurate value for the up-down quark-mixing matrix element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix, V_{ud} . At present, the value of V_{ud} deduced from nuclear beta decay is such that, with standard values [3] of the other elements of the CKM matrix, the unitarity test from the sum of the squares of the elements in the first row fails to meet unity by more than twice the estimated error.

According to CVC, the measured ft -values for Fermi decays closely reflect the value of the weak vector coupling constant, G_V , and are independent of nuclear structure, outside of small correction terms that are of order 1%. Specifically for an isospin-1 multiplet

$$\mathcal{F}t = ft(1 + \delta_R)(1 - \delta_C) = \frac{K}{2G_V'^2}, \quad (1)$$

where f is the statistical rate function, t the partial half-life for the transition, δ_R is the calculated nucleus-dependent radiative correction, δ_C the calculated isospin-breaking correction, and K is a known [1] constant. The effective coupling constant




FIGURE 1. $\mathcal{F}t$ -values for the nine precision data, and the best least-squares one-parameter fit

relates to the primitive one via $G'_v = G_v(1 + \Delta_R^v)^{1/2}$, where Δ_R^v is a calculated nucleus-independent radiative correction. For tests of the CVC hypothesis it is not necessary to consider this correction.

World data on Q -values, lifetimes and branching ratios were thoroughly surveyed [1] in 1989 and updated again [2] for the ENAM95 conference. Since then, there has been a new ^{10}C branching-ratio measurement [4] and a more precise ^{38m}K Q -value determination [5]. We have incorporated both measurements into our data base and extracted the $\mathcal{F}t$ -values plotted in Fig. 1, which also uses the δ_R and δ_C corrections tabulated in our ENAM95 report [2]. It should be noted that those values of δ_C are, in fact, the averages of two independent calculations [6,7]. In a real sense, both experimentally and theoretically, Fig. 1 represents the totality of current world knowledge. The uncertainties shown reflect the experimental uncertainties and an estimate of the *relative* uncertainties in δ_C . There is no statistically significant evidence of inconsistencies in the data ($\chi^2/\nu = 1.1$), thus verifying the expectation of CVC at the level of 3×10^{-4} , the fractional uncertainty quoted on the average $\mathcal{F}t$ -value (3072.3 ± 0.9 s).

In using the average $\mathcal{F}t$ -value to determine V_{ud} and test CKM unitarity it is important to incorporate the ‘systematic’ uncertainty in δ_C that arises from the small systematic differences between the two independent model calculations [6,7]. The result is

$$\mathcal{F}t = 3072.3 \pm 2.0 \text{ s.} \quad (2)$$

With this value, an estimate [8] of the nucleus-independent radiative correction of

$\Delta_R^V = (2.40 \pm 0.08)\%$, and the weak vector coupling constant [3] derived from muon decay, we obtain

$$V_{ud} = 0.9740 \pm 0.0005. \quad (3)$$

The quoted uncertainty is dominated by uncertainties in the theoretical corrections, Δ_R^V and δ_C . On adopting the values [3] of V_{us} and V_{ub} from the Particle Data Group, the sum of squares of the elements in the first row of the CKM matrix,

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9968 \pm 0.0014, \quad (4)$$

differs from unity at the 98% confidence level.

To restore unitarity, the calculated radiative corrections would have to be shifted downwards by 0.3% (*i.e.* as much as one-quarter of their current value), or the calculated Coulomb correction shifted upwards by 0.3% (nearly one-half their value), or some combination of the two. In what follows, we discuss the accuracy of these two corrections and the direction of future research.

RADIATIVE CORRECTIONS

As mentioned, the radiative correction is conveniently divided into terms that are nucleus-dependent, δ_R , and terms that are not, Δ_R^V . These are written

$$\begin{aligned} \delta_R &= \frac{\alpha}{2\pi} [\bar{g}(E_m) + \delta_2 + \delta_3 + 2C_{NS}] \\ \Delta_R^V &= \frac{\alpha}{2\pi} [4 \ln(m_Z/m_p) + \ln(m_p/m_A) + 2C_{\text{Born}}] + \dots, \end{aligned} \quad (5)$$

where the ellipses represent further small terms of order 0.1%. In these equations, E_m is the maximum electron energy in beta decay, m_Z the Z -boson mass, m_A the a_1 -meson mass, and δ_2 and δ_3 the order $Z\alpha^2$ and $Z^2\alpha^3$ contributions. The electron-energy dependent function, $g(E_e, E_m)$ was derived by Sirlin [9]; it is here averaged over the electron spectrum to give $\bar{g}(E_m)$.

Typical values are

$$\delta_R \simeq 0.95 + 0.43 + 0.05 + (\alpha/\pi)C_{NS}\%, \quad (6)$$

where $(\alpha/\pi)C_{NS}$ is of order -0.3% for $T_z = -1$ beta emitters, ^{10}C and ^{14}O , and of order five times smaller for the $T_z = 0$ emitters, ranging from -0.09% to $+0.03\%$ [10]. Thus for $T_z = 0$ emitters $\delta_R \simeq 1.4\%$. If the failure to obtain unitarity in the CKM matrix with V_{ud} from nuclear beta decay is due to the value of δ_R , then δ_R must be reduced to 1.1%. This is not likely. The leading term, 0.95%, involves standard QED and is well verified. The order- $Z\alpha^2$ term, 0.43%, while less secure has been calculated twice [11,12] independently, with results in accord.

For the nucleus-independent term

$$\Delta_{\text{r}}^{\text{v}} = 2.12 - 0.03 + 0.20 + 0.1\% \simeq 2.4\% \quad (7)$$

of which the first term, the leading logarithm, is unambiguous. Again, to achieve unitarity of the CKM matrix, $\Delta_{\text{r}}^{\text{v}}$ would have to be reduced to 2.1%, *i.e.* all terms other than the leading logarithm summing to zero. This also seems unlikely.

COULOMB CORRECTIONS

Because the leading terms in the radiative corrections are well founded, attention has focussed more on the Coulomb correction. Although smaller than the radiative correction, the Coulomb correction is clearly sensitive to nuclear-structure issues. It comes about because Coulomb and charge-dependent nuclear forces destroy isospin symmetry between the initial and final states in superallowed beta-decay. The consequences are twofold: there are different degrees of configuration mixing in the two states, and, because their binding energies are not identical, their radial wave functions differ. Thus we accommodate both effects by writing $\delta_C = \delta_{C1} + \delta_{C2}$. Constraints can be placed on the calculation of δ_{C1} by insisting that the calculation reproduce the coefficients of the isobaric mass multiplet equation. Constraints on δ_{C2} follow by insisting that the asymptotic forms of the proton and neutron radial functions match known separation energies.

Recently Ormand and Brown (OB) [7] have recomputed their Hartree-Fock calculations with new results increasing δ_C over their earlier work [13] but still with values systematically smaller than the Saxon-Woods calculations of Towner, Hardy and Harvey (THH) [6]. Another recent work by Sagawa, van Giai and Suzuki [14] add RPA correlations to a Hartree-Fock calculation; these correlations, in essence, introduce a coupling to the isovector monopole giant resonance. This calculation, however, is not constrained to reproduce known separation energies. Finally a large shell-model calculation has been mounted for the $A = 10$ case by Navrátil, Barrett and Ormand [15]. Both of these two new works [14,15] have produced values of δ_C *smaller* than those used before, *i.e.* worsening rather than helping the unitarity problem.

The typical value of δ_C is of order 0.4%. If the unitarity problem is to be solved by improvements in δ_C , then δ_C has to be raised to around 0.7%. There is no evidence whatsoever for such a shift from recent works.

The δ_C calculations, as pointed out by OB [7], do predict that δ_C should be dramatically larger for nuclei in the fp -shell with $A \geq 62$. This is due to the increasing importance of the $1p$ orbital, which, with its extra node in the radial function compared to the $0f$ orbital, is much more sensitive to Coulomb effects. A similar effect was predicted earlier [6] for $T_z = -1$ nuclei in the middle of the sd -shell where the $1s$ orbital plays an equivalent role. Future experiments will test these predictions.

FUTURE PROSPECTS FOR EXPERIMENT

The nine superallowed transitions surveyed here have been the subject of intense scrutiny for at least the past three decades. All except ^{10}C have the special advantage that the superallowed branch from each is by far the dominant transition in its decay ($> 99\%$). This means that the branching ratio for the superallowed transition can be determined to high precision from relatively imprecise measurements of the other weak transitions, which can simply be subtracted from 100%. Given the quantity of careful measurements already published, are there reasonable prospects for significant improvements in these decays in the near future? Given the uncertainty in the theoretical corrections, perhaps a more important question is whether there is any reason to seek experimental improvements at all.

If we begin by accepting that it is valuable for experiment to be at least a factor of two more precise than theory, then an examination of the world data shows that the Q -values for ^{10}C , ^{14}O , ^{26m}Al and ^{42}Sc , the half-lives of ^{10}C , ^{34}Cl and ^{38m}K , and the branching ratio for ^{10}C can all bear improvement. Such improvements will soon be feasible. The Q -values will reach the required level (and more) as mass measurements with new on-line Penning traps become possible; half-lives will likely yield to measurements with higher statistics as high-intensity beams of separated isotopes are developed for the new radioactive-beam facilities; and, finally, an improved branching-ratio measurement on ^{10}C has already been made with Gammasphere and simply awaits analysis [4].

Qualitative improvements will also come as we increase the number of superallowed emitters accessible to precision studies. The greatest attention recently has been paid to the $T_z = 0$ emitters with $A \geq 62$, since these nuclei are expected to be produced at new radioactive-beam facilities, and their calculated Coulomb corrections, δ_C , are predicted to be large [7,14,16]. They could then provide a valuable test of the accuracy of δ_C calculations. It is likely, though, that the required precision will not be attainable for some time to come. The decays of these nuclei will be of higher energy and each will therefore involve several allowed transitions of significant intensity in addition to the superallowed transition. Branching-ratio measurements will thus be very demanding, particularly with the limited intensities likely to be available initially for these rather exotic nuclei. Lifetime measurements will be similarly constrained by statistics.

More accessible in the short term will be the $T_z = -1$ superallowed emitters with $18 \leq A \leq 38$. There is good reason to explore them. For example, the calculated value [6] of δ_C for ^{30}S decay, though smaller than the δ_C 's expected for the heavier nuclei, is actually 1.2% – about a factor of two larger than for any other case currently known – while ^{22}Mg has a very low value of 0.35%. If such large differences are confirmed by the measured ft -values, then it will do much to increase our confidence in the calculated Coulomb corrections. To be sure, these decays will provide a challenge, particularly in the measurement of their branching ratios, but the required precision should be achievable with isotope-separated beams that are currently available. In fact, such experiments are already in their early stages at

the Texas A&M cyclotron.

CONCLUSIONS

The current world data on superallowed $0^+ \rightarrow 0^+$ beta decays lead to a self-consistent set of $\mathcal{F}t$ -values that agree with CVC but differ provocatively, though not yet definitively, from the expectation of CKM unitarity. There are no evident defects in the calculated radiative and Coulomb corrections that could remove the problem, so, if any progress is to be made in firmly establishing (or eliminating) the discrepancy with unitarity, additional experiments are required. We have indicated what some relevant nuclear experiments might be.

Clearly, there is strong motivation to pursue them since, if firmly established, a discrepancy with unitarity would indicate the need for an extension of the three-generation Standard Model.

REFERENCES

1. J.C. Hardy, I.S. Towner, V.T. Koslowsky, E. Hagberg and H. Schmeing, Nucl. Phys. **A509**, 429 (1990)
2. I.S. Towner, E. Hagberg, J.C. Hardy, V. Koslowsky and G. Savard, *Proceedings of the Int. Conf. on Exotic Nuclei and Atomic Masses, Arles, France, June 1995*, eds. M. de Saint Simon and O. Sorlin (Edition Frontières, Gif-sur-Yvette, France, 1995) pp.711-721
3. C. Caso *et al.*, The European Physical Journal **C3**, 1 (1998)
4. B.K. Fujikawa *et al.*, APS e-print 1998may29_004
5. P.H. Barker, P.D. Harty and N.S. Bowden, private communication
6. I.S. Towner, J.C. Hardy and M. Harvey, Nucl. Phys. **A284**, 269 (1977)
7. W.E. Ormand and B.A. Brown, Phys. Rev. **C52**, 2455 (1995)
8. A. Sirlin, in *Precision Tests of the Standard Electroweak Model*, ed. P. Langacker (World-Scientific, Singapore, 1994)
9. A. Sirlin, Phys. Rev. **164**, 1767 (1967)
10. I.S. Towner, Phys. Lett. **B333**, 13 (1994)
11. A. Sirlin, Phys. Rev. **D35**, 3423 (1987); A. Sirlin and R. Zucchini, Phys. Rev. Lett. **57**, 1994 (1986)
12. W. Jaus and G. Rasche, Phys. Rev. **D35**, 3420 (1987)
13. W.E. Ormand and B.A. Brown, Phys. Rev. Lett. **62**, 866 (1989); Nucl. Phys. **A440**, 274 (1985)
14. H. Sagawa, N. van Giai and T. Suzuki, Phys. Rev. **C53**, 2163 (1996)
15. P. Navrátil, B.R. Barrett and W.E. Ormand, Phys. Rev. **C56**, 2542 (1997)
16. J.C. Hardy, in *Proceedings of the Workshop on the Production and Use of Intense Radioactive Beams at the Isospin Laboratory*, ed. J.D. Garrett (Joint Institute for Heavy Ion Research, Oak Ridge, Tennessee: Conf. 9210121) pg. 51